


MAS331: Metric Spaces 2016–17
Solutions to Week 2 Problems on Chapter 1


5. Let $r > 0$ and let $a = (a_1, \dots, a_n), b = (b_1, \dots, b_n) \in \mathbb{R}^n$. Suppose that $b \in B_1(a, r)$. Then $d_1(b, a) < r$. But $d_1(a, b) \geq d_2(a, b)$ by 3(a) so $d_2(a, b) \leq d_1(b, a) < r$ and hence $b \in B_2(a, r)$. Thus $B_1(a, r) \subseteq B_2(a, r)$.

Now suppose that $b \in B_2(a, r)$. Then $d_2(b, a) < r$. But $d_2(a, b) \geq d_\infty(a, b)$ by 3(a) so $d_\infty(a, b) \leq d_2(b, a) < r$ and hence $b \in B_\infty(a, r)$. Thus $B_2(a, r) \subseteq B_\infty(a, r)$.




$$B_1(a, r) \subseteq B_2(a, r) \subseteq B_\infty(a, r)$$

Next, suppose that $b \in B_\infty(a, \frac{r}{n})$. Then $d_\infty(b, a) < \frac{r}{n}$. Using 3(b), $d_1(a, b) \leq n d_\infty(a, b) < r$ so $b \in B_1(a, r)$. Thus $B_\infty(a, \frac{r}{n}) \subseteq B_1(a, r)$.



$$B_\infty(a, \frac{r}{n}) \subseteq B_1(a, r) \subseteq B_\infty(a, r)$$

Finally, suppose that $b \in B_\infty(a, \frac{r}{\sqrt{n}})$. Then $d_\infty(b, a) < \frac{r}{\sqrt{n}}$. Using 3(c), $d_2(a, b) \leq \sqrt{n} d_\infty(a, b) < r$ so $b \in B_2(a, r)$. Thus $B_\infty(a, \frac{r}{\sqrt{n}}) \subseteq B_2(a, r)$.




$$B_\infty(a, \frac{r}{\sqrt{n}}) \subseteq B_2(a, r) \subseteq B_\infty(a, r)$$

$$B_\infty(a, \frac{r}{n}) \subseteq B_1(a, r) \text{ and } B_\infty(a, \frac{r}{\sqrt{n}}) \subseteq B_2(a, r).$$

Here, as in the notes, the subscripts indicate which metric we are using.

When $n > 1$, all the inclusions \subseteq are strict inclusions.

6. Here's a picture of the graphs between 0 and 5:



To compute $d_\infty(f_1, f_2)$, we need to find the maximum value for $|f(x) - g(x)|$ with $x \in I$. The only possible x values to consider are stationary points of $f - g$ and the two endpoints of the interval I .

At $x = 0$, $|f(x) - g(x)| = 6$; at $x = 5$, $|f(x) - g(x)| = 4$. Next, let's work out the stationary points of $f - g$. The difference $f(x) - g(x)$ is $x^2 - 7x + 6$, so its derivative is $2x - 7$; thus there is only one stationary point, at $x = \frac{7}{2}$, and the graphs are a distance of $\frac{25}{4}$ apart there. Since $\frac{25}{4} > 6$, this is the required answer. So

$$d_\infty(x^2 - 4x, 3x - 6) = \frac{25}{4}.$$

To compute $d_1(f_1, f_2)$, we calculate the area between the two graphs. Now the graphs meet when $x^2 - 4x = 3x - 6$, which occurs at $x = 1$ (and again outside the interval $[0, 5]$). We have $x^2 - 4x \geq 3x - 6$ for $0 \leq x \leq 1$, and $x^2 - 4x \leq 3x - 6$ for $1 \leq x \leq 5$. So

$$\begin{aligned} d_1(x^2 - 4x, 3x - 6) &= \int_0^5 |(x^2 - 4x) - (3x - 6)| dx \\ &= \int_0^1 (x^2 - 4x) - (3x - 6) dx + \int_1^5 (3x - 6) - (x^2 - 4x) dx \\ &= \left[\frac{x^3}{3} - \frac{7x^2}{2} + 6x \right]_0^1 + \left[\frac{7x^2}{2} - \frac{x^3}{3} - 6x \right]_1^5 \\ &= \left(\frac{17}{6} - 0 \right) + \left(\frac{95}{6} - \left(-\frac{17}{6} \right) \right) \\ &= \frac{129}{6} = \frac{43}{2}. \end{aligned}$$

- 7.

$$B[f, 1] = \{g \in C[0, 1] \mid d_\infty(f, g) \leq 1\},$$

and it follows from the definitions that

$$d_\infty(f, g) = \sup\{|g(x) - f(x)| : x \in [0, 1]\} = \sup\{|g(x)| : x \in [0, 1]\}$$

so that $d_\infty(f, g) \leq 1$ if and only if $|g(x)| \leq 1$ for all $x \in [0, 1]$. This means that the closed ball is the set of functions whose graph lies between the lines $y = 1$ and $y = -1$.

8.

$$d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{otherwise} \end{cases}$$

It is clear from this definition that d satisfies M1 and M2.

It remains to verify the triangle inequality M3. Let $x, y, z \in X$. We want to show that $d(x, y) \leq d(x, z) + d(z, y)$. If $d(x, y) = 0$ then the inequality obviously holds. Therefore we assume that $d(x, y) = 1$ so that $x \neq y$. It follows that z must be different from one of x or y so that $d(x, z) = 1$ or $d(z, y) = 1$ or both. Hence

$$d(x, z) + d(z, y) \geq 1 = d(x, y).$$

9. Let $f, g \in C[a, b]$. Then, since $|f(x) - g(x)| = |g(x) - f(x)|$ for all $x \in [a, b]$, we have

$$\begin{aligned} d_1(f, g) &= \int_a^b |f(x) - g(x)| dx \\ &= \int_a^b |g(x) - f(x)| dx \\ &= d_1(g, f), \end{aligned}$$

so that d_1 satisfies axiom M2.

Now let $h \in C[a, b]$. Since

$$|f(x) - h(x)| = |(f(x) - g(x)) + (g(x) - h(x))| \leq |f(x) - g(x)| + |g(x) - h(x)|$$

for all $x \in [a, b]$, we have

$$\begin{aligned} d_1(f, h) &= \int_a^b |f(x) - h(x)| dx \\ &\leq \int_a^b |f(x) - g(x)| + |g(x) - h(x)| dx \\ &= \int_a^b |f(x) - g(x)| dx + \int_a^b |g(x) - h(x)| dx \\ &= d_1(f, g) + d_1(g, h), \end{aligned}$$

which proves M3.

10. (a) $(4, 2)$, $(2, 1)$ and the origin are collinear so $d((4, 2), (2, 1)) = d_2((4, 2), (2, 1)) = \sqrt{5}$. $(4, 2)$, $(-2, 3)$ and the origin are not collinear so we have to go via the origin and


$$d((4, 2), (-2, 3)) = d_2((4, 2), (0, 0)) + d_2((-2, 3), (0, 0)) = 2\sqrt{5} + \sqrt{13}.$$

- (b) $(0, -1)$ is at distance 1 from $(0, 0)$ so the only points within $\frac{1}{2}$ of $(0, -1)$ are on the line through $(-1, 0)$ and $(0, 0)$ (the y -axis) so

$$B((0, -1), \frac{1}{2}) = \{(0, y) : -0.5 > y > -1.5\}.$$

There are two types of points in $B((0, -1), 2)$. There are those that are collinear with $(-1, 0)$ and $(0, 0)$ at distance < 2 from $(-1, 0)$. That gives $\{(0, y) : 1 > y > -3\}$. There are those that we reach via the $(0, 0)$ which must be < 1 from $(0, 0)$. That gives $\{(x, y) : x^2 + y^2 < 1\}$. So

$$B((0, -1), 2) = \{(x, y) : x^2 + y^2 < 1 \text{ or } (x = 0 \text{ and } -3 < y < 1)\}.$$



In the right-hand diagram $(-1, 0)$ is included but all other points on the unit circle are excluded. In the left-hand diagram, both endpoints, $(0, -0.5)$ and $(0, -1.5)$, are excluded.

11. First do the case $d(x, y) = \sqrt{|x - y|}$. Take $x, y, z \in \mathbb{R}$. For M1, first $d(x, x) = \sqrt{|x - x|} = 0$, and then if $\sqrt{|x - y|} = d(x, y) = 0$ we find $|x - y| = (\sqrt{|x - y|})^2 = 0$, so that $x = y$.

For M2, we simply note that $d(y, x) = \sqrt{|y - x|} = \sqrt{|x - y|} = d(x, y)$.

For M3, we want to know if

$$d(x, y) = \sqrt{|x - y|} \leq d(x, z) + d(z, y) = \sqrt{|x - z|} + \sqrt{|z - y|}.$$

If we square both sides, we end up asking if the inequality

$$|x - y| \leq |x - z| + |z - y| + 2\sqrt{|x - z||z - y|}$$

holds. The answer clearly is yes since we already have $|x - y| \leq |x - z| + |z - y|$.

In the second case, if we take $x = 1$, $y = -1$ and $z = 0$ we find that $d(x, y) = (1 - (-1))^2 = 4$, but $d(x, z) + d(z, y) = 1^2 + (-1)^2 = 2$. Thus M3 is not satisfied and this d is not a metric. (Here M1 and M2 are satisfied.)

12. For M1, we first check that $d(x, y) \geq 0$ for all $x, y \in X$: as d_1 and d_2 satisfy M1,

$$d(x, y) = d_1(x, y) + d_2(x, y) \geq 0.$$

Also, as d_1 and d_2 satisfy M1, $d(x, x) = d_1(x, x) + d_2(x, x) = 0 + 0 = 0$. And if $d(x, y) = 0$ then, as $d_1(x, y) \geq 0$ and $d_2(x, y) \geq 0$, we must have $d_1(x, y) = d_2(x, y) = 0$ and so $x = y$.

For M2, d_1 and d_2 satisfy M2 so

$$d(x, y) = d_1(x, y) + d_2(x, y) = d_1(y, x) + d_2(y, x) = d(y, x).$$

For M3, d_1 and d_2 satisfy M3 so, for $x, y, z \in X$,

$$\begin{aligned} d(x, z) &= d_1(x, z) + d_2(x, z) \\ &\leq (d_1(x, y) + d_1(y, z)) + (d_2(x, y) + d_2(y, z)) \\ &= (d_1(x, y) + d_2(x, y)) + (d_1(y, z) + d_2(y, z)) \\ &= d(x, y) + d(y, z). \end{aligned}$$

Thus d satisfies M1,2,3 and d is a metric on X .

13. $d_1((0, 0), (1, 0)) = 1 = d_2((0, 0), (1, 0))$ so $d((0, 0), (1, 0)) = 0$. As $(0, 0) \neq (1, 0)$, d fails axiom M1 and is therefore not a metric.
14. (a) Let $n = 2$. Then one of $|x_1 - y_1|, |x_2 - y_2|$ is $\max(|x_i - y_i|)$ and the other is $\min\{|x_i - y_i|\}$. This includes the case where $|x_1 - y_1| = |x_2 - y_2|$ and $\max(|x_i - y_i|) = \min(|x_i - y_i|)$. Hence

$$\max(|x_i - y_i|) + \min(|x_i - y_i|) = |x_1 - y_1| + |x_2 - y_2|,$$

that is, $e(x, y) = d_1(x, y)$.

- (b) Let $x = (0, 0, 0)$, $y = (1, 2, 3)$ and $z = (2, 2, 4)$.

Then $e(x, y) = 3 + 1 = 4$, $e(y, z) = 1 + 0 = 1$. But $e(x, z) = 4 + 2 = 6 > e(x, y) + e(y, z)$ so axiom M3 fails and e is not a metric.