

Infinitely Divisible Central Probability Measures on Compact Lie Groups - Regularity, Semigroups and Transition Kernels

David Applebaum,
Department of Probability and Statistics,
University of Sheffield,
Hicks Building, Hounsfield Road,
Sheffield, England, S3 7RH

e-mail: D.Applebaum@sheffield.ac.uk

Abstract

We introduce a class of central symmetric infinitely divisible probability measures on compact Lie groups by lifting the characteristic exponent from the real line via the Casimir operator. The class includes Gauss, Laplace and stable-type measures. We find conditions for such a measure to have a smooth density and give examples. The Hunt semigroup and generator of convolution semigroups of measures are represented as pseudo-differential operators. For sufficiently regular convolution semigroups, the transition kernel has a tractable Fourier expansion and the density at the neutral element may be expressed as the trace of the Hunt semigroup. We compute the short time asymptotics of the density at the neutral element for the Cauchy distribution on the d -torus, on $SU(2)$ and on $SO(3)$, where we find markedly different behaviour than is the case for the usual heat kernel.

Key words and phrases infinite divisibility, central measure, compact Lie group, Casimir operator, Sobolev space, convolution semigroup, Hunt semigroup, pseudo-differential operator, symbol, transition density

AMS 2000 subject classification 60B15, 60G51, 47D07, 43A05, 35K08

1 Introduction

The heat kernel on a compact Riemannian manifold has been the subject of extensive investigations by analysts, geometers and probabilists. One reason for this is that its small and large time asymptotic expansions contain important topological and geometric information (see e.g. [32]). Another reason is that it is the transition density of manifold-valued Brownian motion which is a stochastic process of intrinsic interest (see e.g. [10]). If the manifold is a Lie group then the heat kernel is naturally related to Dedekind's eta function via Macdonald's identities (see [12]). In this paper we will mainly be concerned with compact Lie groups. In this case the heat kernel is naturally associated to a vaguely (or equivalently, weakly) continuous convolution semigroup of probability measures which we'll refer to as the "heat semigroup" in the sequel.

The study of the entire class of such convolution semigroups has had a long development (see e.g. [20], [19], [5]). From a probabilistic point of view they correspond to Lévy processes, i.e. stochastic processes with stationary and independent increments. Compared to Brownian motion which has continuous sample paths (with probability one) the paths of the generic Lévy process are only right continuous and have jump discontinuities of arbitrary size occurring at random times.

The purpose of this paper is to study a class of convolution semigroups which on the one hand, are sufficiently close in structure to the heat semigroup to enable us to do some interesting analysis and on the other hand, are sufficiently broad as to display all the interesting features that one finds with a generic Lévy process. The first observation is that measures comprising the heat semigroup are central and so we focus on this class. Its worth pointing out that that central measures as a class have also received some attention from analysts (see e.g. [31], [16]). Secondly we remark that if $(\mu_t, t \geq 0)$ is the heat semigroup then its non-commutative Fourier transform (see [17], [36] for background on this concept) takes the form $\widehat{\mu}_t(\pi) = e^{-\frac{t}{2}\kappa_\pi} I_\pi$ at the irreducible representation π where $-\kappa_\pi I_\pi$ is the Casimir operator. But $u \rightarrow \frac{u^2}{2}$ is the negative-definite function on the real line associated with the standard Gaussian measure. The generalisation that we make here is to consider a class of semigroups that are given by the prescription $\widehat{\mu}_t(\pi) = e^{-t\eta(\kappa_\pi^{\frac{1}{2}})} I_\pi$ where η is a symmetric negative definite function. Other examples of measures subsumed within this class include the Laplace distribution on a Lie group, which has been utilised in recent statistical work on the problem of deconvolution (see e.g. [26], [24]) and analogues of stable laws. Indeed any semigroup of probability measures that is obtained by subordinating the heat

semigroup belongs to this class. We study these measures both from the static perspective, where the emphasis is on a single infinitely divisible measure, and the dynamic perspective where we focus on an entire semigroup.

The organisation of this paper is as follows. In section 2, we study central probability measures, introduce our main class and examine some examples. In section 3 we use Sobolev spaces to find conditions on our induced measures which enable them to have a smooth density. In section 4 we turn our attention to convolution semigroups and the associated semigroup of operators (the *Hunt semigroup*) on the L^2 space of normalised Haar measure. When G is a Euclidean space it is known (see [22], Chapter 3 of [4]) that these operators, and their generators, can be realised as pseudo-differential operators. Using Peter-Weyl theory, Ruzhansky and Turunen [33] have developed an intrinsic theory of pseudo-differential operators on compact groups. We adapt this theory to our needs and show that the Hunt semigroup and its generator are pseudo-differential operators in the sense of Ruzhansky and Turunen. This part of the work is carried out in full generality. In the case of our induced class, we show that the generator has the same Sobolev regularity as the Laplacian. Although these results have an analytic flavour, they are important for probabilists as they indicate a route to investigate general classes of Feller-Markov processes on compact Lie groups using the symbol of the generator as the key tool (see [23] for an account of this theory in the case where G is Euclidean space.)

In section 5, we investigate the transition kernel for convolution semigroups of central measures. We remark that the first investigation of densities for such measures (under a hypo-ellipticity condition that we don't require here) were made by Liao in [28] (see also Theorem 4.4 in [29] p.96.) A necessary and sufficient condition for the semigroup to be trace-class for any positive time is that the corresponding probability measure has a square-integrable density [5]. We compute the trace in both the main L^2 -space and the subspace of central functions. In the former case, the co-ordinate functions form a complete set of eigenvectors for the Hunt semigroup. Comparing the traces in these two spaces leads to an interesting inequality for transition kernels which appears to be new even in the heat kernel case. Finally in section 6, we study the small time asymptotics of the transition kernel in the case of the Cauchy distribution on the d -torus, on $SU(2)$ and on $SO(3)$ and show that it blows up much faster than the heat kernel.

Acknowledgements. I would like to thank Natesh Pillai for many useful discussions and Ming Liao and Michael Ruzhansky for helpful comments. Both René Schilling and Rodrigo Bañuelos provided very valuable remarks after I presented a talk based on part of this paper at the 2010 Lévy processes

conference in Dresden. Last, but not least, I am grateful to the referee for his careful reading and a number of helpful observations.

2 Infinite Divisibility of Central Measures

Let G be a compact group with neutral element e and let $\mathcal{M}(G)$ be the set of all probability measures defined on $(G, \mathcal{B}(G))$ where $\mathcal{B}(G)$ is the Borel σ -algebra of G . We say that $\mu \in \mathcal{M}(G)$ is *central* (or *conjugate-invariant*) if $\mu(\sigma A \sigma^{-1}) = \mu(A)$ for all $\sigma \in G, A \in \mathcal{B}(G)$ and μ is said to be *symmetric* if $\mu(A^{-1}) = \mu(A)$ for all $A \in \mathcal{B}(G)$. Let $\mathcal{M}_c(G)$ ($\mathcal{M}_s(G)$) be the subsets of $\mathcal{M}(G)$ comprising central (symmetric) measures (respectively) and define $\mathcal{M}_{c,s}(G) := \mathcal{M}_c(G) \cap \mathcal{M}_s(G)$. Normalised Haar measure on G will always be denoted $d\sigma$ when integrating functions of $\sigma \in G$.

Let \widehat{G} be the set of all equivalence classes of irreducible representations of G . We will, without further comment, frequently identify equivalence classes with a particular representative element when there is no loss of generality. The trivial representation will always be denoted by δ . Each $\pi \in \widehat{G}$ acts as a $d_\pi \times d_\pi$ unitary matrix on a complex linear space V_π having dimension d_π . We define the Fourier transform of each $\mu \in \mathcal{M}(G)$ to be the Bochner integral

$$\widehat{\mu}(\pi) = \int_G \pi(\sigma) \mu(d\sigma),$$

where $\pi \in \widehat{G}$. We will frequently use the well-known and easily verified fact that

$$\widehat{\mu * \nu}(\pi) = \widehat{\mu}(\pi) \widehat{\nu}(\pi),$$

for all $\mu, \nu \in \mathcal{M}(G), \pi \in \widehat{G}$, where $*$ denotes convolution of measures.

Suppose we are given $\mu \in \mathcal{M}(G)$. It is shown in [34] that $\mu \in \mathcal{M}_c(G)$ if and only if for each $\pi \in \widehat{G}$ there exists $c_\pi \in \mathbb{C}$ such that

$$\widehat{\mu}(\pi) = c_\pi I_\pi, \tag{2.1}$$

where I_π is the identity matrix acting on V_π . Indeed this is a straightforward consequence of Schur's lemma. Moreover one has the formula,

$$c_\pi = \frac{1}{d_\pi} \int_G \chi_\pi(\sigma) \mu(d\sigma), \tag{2.2}$$

where $\chi_\pi(\cdot) := \text{tr}(\pi(\cdot))$ is the group character.

It is well-known (and easily verified) that $\mu \in \mathcal{M}_s(G)$ if and only if $\widehat{\mu}(\pi)$ is self-adjoint for all $\pi \in \widehat{G}$. Consequently $\mu \in \mathcal{M}_{cs}(G)$ if and only if $\widehat{\mu}(\pi) = c_\pi I_\pi$ with $c_\pi \in \mathbb{R}$ for all $\pi \in \widehat{G}$.

A probability measure μ is *infinitely divisible* if for each $n \in \mathbb{N}$ there exists $\nu_n \in \mathcal{M}(G)$ such that $\nu_n^{*n} = \mu$. In this case we write $\mu_{\frac{1}{n}} := \nu_n$.

Proposition 2.1 *If G is a compact Lie group and $\mu \in M_{cs}(G)$ is infinitely divisible then for each $\pi \in \widehat{G}$ there exists $\alpha_\pi \leq 0$ such that $\widehat{\mu}(\pi) = e^{\alpha_\pi} I_\pi$.*

Proof. By the results on pp.220-1 of [19], μ may be embedded as μ_1 into a vaguely continuous convolution semigroup of probability measures $(\mu_t, t \geq 0)$ where μ_0 is normalised Haar measure on a closed subgroup H of G . It follows (see [1], [30]) that for each $\pi \in \widehat{G}$, $(\widehat{\mu}_t(\pi), t \geq 0)$ is a strongly continuous contraction semigroup of matrices acting on V_π and so we may write $\widehat{\mu}_t(\pi) = \widehat{\mu}_0(\pi) e^{tA_\pi}$ for all $t \geq 0$ where A_π is a $d_\pi \times d_\pi$ matrix. Now since $\mu_1 \in M_{cs}(G)$ there exists $\lambda_\pi \in \mathbb{R}$ such that

$$\widehat{\mu}_1(\pi) = \widehat{\mu}_0(\pi) e^{A_\pi} = \lambda_\pi I_\pi \dots (*)$$

If $\lambda_\pi = 0$, the required result holds with $\alpha_\pi = -\infty$ so assume that $\lambda_\pi \neq 0$. Since $\mu_1 = \mu_1 * \mu_0$ we have

$$\widehat{\mu}_0(\pi) e^{A_\pi} \widehat{\mu}_0(\pi) = \lambda_\pi I_\pi.$$

On the other hand, post-multiplying both sides of (*) by $\widehat{\mu}_0(\pi)$ yields

$$\widehat{\mu}_0(\pi) e^{A_\pi} \widehat{\mu}_0(\pi) = \lambda_\pi \widehat{\mu}_0(\pi).$$

It follows that $\widehat{\mu}_0(\pi) = I_\pi$ and hence $H = \{e\}$. We then have $A_\pi = \alpha_\pi I_\pi$ where $\alpha_\pi \in \mathbb{R}$ and $\lambda_\pi = e^{\alpha_\pi}$. But $\widehat{\mu}_1(\pi)$ is a contraction on V_π and hence $\alpha_\pi \leq 0$. \square

Example. *The Compound Poisson Distribution.* Consider the probability measure $\mu_{\lambda, \gamma}$ where γ is a given probability measure on G and $\lambda > 0$. This is defined by

$$\mu_{\lambda, \gamma} := e^{-\lambda} \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} \gamma^{*n}.$$

It is well known (see e.g. [34]) that for all $\pi \in \widehat{G}$,

$$\widehat{\mu}_{\lambda, \gamma}(\pi) = \exp \{ \lambda (\widehat{\gamma}(\pi) - I_\pi) \}.$$

Proposition 2.2 1. The measure $\mu_{\lambda,\gamma}$ is central if and only if γ is.

2. The measure $\mu_{\lambda,\gamma}$ is symmetric if and only if γ is.

Proof.

1. The if part is straightforward and is established in Proposition 4 of [34].
Conversely if $\mu_{\lambda,\gamma}$ is central then for all $g \in G, \pi \in \widehat{G}$

$$\pi(g)\widehat{\mu_{\lambda,\gamma}}(\pi)\pi(g^{-1}) = \widehat{\mu_{\lambda,\gamma}}(\pi),$$

and so

$$\widehat{\mu_{\lambda,\gamma}}(\pi) = \exp \{ \lambda(\pi(g)\widehat{\gamma}(\pi)\pi(g^{-1}) - I_\pi) \}.$$

Now by uniqueness of Fourier transforms and injectivity of the exponential map on matrices, we have

$$\pi(g)\widehat{\gamma}(\pi)\pi(g^{-1}) = \widehat{\gamma}(\pi)$$

for all $g \in G, \pi \in \widehat{G}$ and the result follows.

2. This result is proved similarly using the fact that a probability measure is symmetric if and only if its Fourier transform comprises self-adjoint matrices.

□

It follows that a central probability measure μ is a compound Poisson distribution if and only if there exists $\lambda > 0$ and a central probability measure γ with $\widehat{\gamma}(\pi) = b_\pi I_\pi$ for all $\pi \in \widehat{G}$ such that

$$\widehat{\mu}(\pi) = \exp \{ \lambda(b_\pi - 1)I_\pi \}. \quad (2.3)$$

We now introduce a class of central symmetric measures which are key to this paper. For this part we assume that G is a compact Lie group. Let ρ be a symmetric infinitely divisible probability measure on \mathbb{R} . Then we have the Lévy-Khintchine formula

$$\int_{\mathbb{R}} e^{iux} \rho(dx) = e^{-\eta(u)},$$

where

$$\eta(u) = \frac{1}{2}\sigma^2 u^2 + \int_{\mathbb{R}-\{0\}} (1 - \cos(uy))\nu(dy), \quad (2.4)$$

where $\sigma \geq 0$ and ν is a symmetric Lévy measure on $\mathbb{R} - \{0\}$, i.e. a σ -finite Borel symmetric measure for which $\int_{\mathbb{R}-\{0\}} \min\{1, |x|^2\} \nu(dx) < \infty$ (see e.g. [6].)

For each $\pi \in \widehat{G}$ let K_π be the Casimir operator acting in V_π . Then $K_\pi = -\kappa_\pi I_\pi$ where $\kappa_\pi \geq 0$ with $\kappa_\pi = 0$ if and only if $\pi = \delta$. If μ is a probability measure on G for which

$$\widehat{\mu}(\pi) = e^{-\eta(\kappa_\pi^{\frac{1}{2}})} I_\pi, \quad (2.5)$$

we say that μ is a central symmetric probability measure on G induced by an infinitely divisible probability measure on \mathbb{R} and we write $\mu \in \text{CID}_{\mathbb{R}}(G)$.

The following two examples have been applied to statistical inference on groups (see e.g. [24], [26].)

Example 1. *Gaussian measure.* Here we take $\nu = 0$ and so $c_\pi = \exp\{-\frac{1}{2}\sigma^2\kappa_\pi\}$. Gaussian measure is embeddable into the Brownian motion or heat semigroup of measures for which $\widehat{\mu}_t(\pi) = \exp\{-\frac{t}{2}\sigma^2\kappa_\pi\}$ for $t \geq 0$ which has been extensively studied by both analysts and probabilists.

Example 2. *The Laplace distribution on G .* Here we take $\sigma = 0, \nu(dx) = \frac{\exp\{-\frac{|x|}{\beta}\}}{|x|} dx$ (with $\beta > 0$) and $c_\pi = (1 + \beta^2\kappa_\pi)^{-1}$ (see [35], p.98 for a discussion of the underlying distribution on \mathbb{R}).

Now consider a central symmetric compound Poisson distribution $\mu_{\lambda,\gamma}$. We consider conditions under which $\mu_{\lambda,\gamma} \in \text{CID}_{\mathbb{R}}(G)$. First take $\sigma = 0$ and ν to be a finite symmetric measure in (2.4) and rewrite

$$\eta(u) = \lambda \int_{\mathbb{R}-\{0\}} (1 - \cos(uy)) \tilde{\nu}(dy),$$

where $\lambda := \nu(\mathbb{R} - \{0\})$ and $\tilde{\nu}(\cdot) := \frac{1}{\lambda} \nu(\cdot)$. For $\mu_{\lambda,\gamma} \in \text{CID}_{\mathbb{R}}(G)$ with this value of λ we require that $b_\pi = g(\kappa_\pi^{\frac{1}{2}})$ in (2.3) where $g(u) = \int_{\mathbb{R}} \cos(ux) \tilde{\nu}(dx)$. For example if we take ν to be a constant multiple of a centred Gaussian measure with variance σ^2 on \mathbb{R} then $b_\pi = \exp\{-\frac{1}{2}\sigma^2\kappa_\pi\}$.

We now consider an important subclass of measures in $\text{CID}_{\mathbb{R}}(G)$. Let $(\rho_t^f, t \geq 0)$ be the law of a subordinator with associated Bernstein function $f : (0, \infty) \rightarrow [0, \infty)$ so that $(\rho_t^f, t \geq 0)$ is a vaguely continuous convolution semigroup of probability measures on $[0, \infty)$ and for each $t \geq 0, u > 0$,

$$\int_0^\infty e^{-us} \rho_t^f(ds) = e^{-tf(u)}, \quad (2.6)$$

and f has the generic form

$$f(u) = au + \int_{(0,\infty)} (1 - e^{-uy})\lambda(dy),$$

where $a \geq 0$ and $\int_{(0,\infty)} \min\{1, y\}\lambda(dy) < \infty$ (see e.g. [35], section 30 and [4], section 1.3.2 for details.) It is straightforward to verify that if $(\mu_t, t \geq 0)$ is a vaguely continuous convolution semigroups of measures on G and $(\rho_t^f, t \geq 0)$ is a subordinator as above then we get another vaguely continuous convolution semigroups of measures on G which we denote $(\mu_t^f, t \geq 0)$ via the vague integral

$$\mu_t^f(A) = \int_0^\infty \mu_s(A)\rho_t^f(ds),$$

for $A \in \mathcal{B}(G)$. Now let $(\mu_t, t \geq 0)$ be the Brownian motion semigroup with $\sigma = \sqrt{2}$. Then for each $\pi \in \widehat{G}, t \geq 0$ we have

$$\begin{aligned} \widehat{\mu}_t^f(\pi) &= \int_0^\infty \int_G \pi(\sigma)\mu_s(d\sigma)\rho_t^f(ds) \\ &= \left(\int_0^\infty e^{-s\kappa_\pi}\rho_t^f(ds) \right) I_\pi \\ &= e^{-tf(\kappa_\pi)} I_\pi, \end{aligned}$$

and so $\mu_1^f \in \text{CID}_{\mathbb{R}}(G)$ with $\eta(\kappa_\pi^{\frac{1}{2}}) = f(\kappa_\pi)$.

Note that the Laplace distribution (as described above) is obtained in this way with $f(u) = \log(1 + \beta^2 u)$. It is worth pointing out that it also arises as $\beta^{-2}V^{\beta^{-2}}$ where for $c > 0, V^c$ is the potential measure of the Brownian motion semigroup defined by the vague integral $V^c(\cdot) = \int_0^\infty e^{-ct}\mu_t(\cdot)dt$ (see [35], pp. 203-5 for the case in \mathbb{R}^d .)

Other examples of measures in $\text{CID}_{\mathbb{R}}(G)$ which are obtained by subordination include stable-type distributions where $\sigma = 0$ in (2.4) and $\nu(dx) = \frac{b^\alpha}{|x|^{1+\alpha}}dx$ where $b > 0$ and $0 < \alpha < 2$. In this case we have $f(u) = b^\alpha u^{\frac{\alpha}{2}}$ and $c_\pi = \exp\left\{-b^\alpha \kappa_\pi^{\frac{\alpha}{2}}\right\}$. We may also consider the relativistic Schrödinger distribution for $m > 0$ where $f(u) = \sqrt{u + m^2} - m$ and $c_\pi = e^{-(\sqrt{m^2 + \kappa_\pi} - m)}$. It again has $\sigma = 0$ in (2.4). The precise form of ν is complicated and as we don't require it here we refer the reader to [21].

It is an interesting problem to determine the class of all η in (2.4) which give rise to a probability measure on G of the form (2.5).

3 Regularity of Densities

In this section we will assume that G is a compact semi-simple Lie group having Lie algebra \mathfrak{g} . We say that $\mu \in \mathcal{M}(G)$ has a density $k \in L^1(G, \mathbb{R})$ if μ is absolutely continuous with respect to normalised Haar measure on G .

We then define k to be the Radon-Nikodým derivative $\frac{d\mu}{d\sigma}$.

If a density k exists for $\mu \in \mathcal{M}_c(G)$ with $\widehat{\mu}(\pi) = c_\pi I_\pi$ and $k \in L^2(G, \mathbb{R})$ then it has the form

$$k(\sigma) = \sum_{\pi \in \widehat{G}} d_\pi \overline{c_\pi} \chi_\pi(\sigma), \quad (3.7)$$

for almost all $\sigma \in G$ (see equation (3.4) in [3].)

Before we investigate densities in greater detail we need some preliminaries.

3.1 Dominant weights

Fix a maximal torus \mathbb{T} in G . Let \mathbb{T} be its Lie algebra and \mathbb{T}^* be the dual vector space to \mathbb{T} . Let P be the lattice of weights in \mathbb{T}^* and $D \subset \mathbb{T}^*$ be the dominant chamber. The celebrated theorem of the highest weight asserts that there is a one-to-one correspondence between elements of \widehat{G} and the highest weights which are precisely the members of $P \cap D$. For details see e.g. Chapters 8 and 9 in [14]. In the following the inner product $\langle \cdot, \cdot \rangle$ and norm $|\cdot|$ on \mathbb{T}^* are that induced by the Killing form via duality.

Let λ_π be the dominant weight for the representation π . Then we know from Sugira [37] (p. 39, equation (1.17)) that

$$d_\pi \leq N |\lambda_\pi|^m \quad (3.8)$$

where N is a universal constant and

$$m = \frac{1}{2}(\dim(G) - r) \quad (3.9)$$

where r is the rank of G , i.e. the dimension of any maximal torus. It is also well known that (see e.g. [37], Lemma 1.1 or [25], Proposition 5.28)

$$\kappa_\pi = |\lambda_\pi + \rho|^2 - |\rho|^2 = \langle \lambda_\pi, \lambda_\pi + 2\rho \rangle \quad (3.10)$$

where ρ is half the sum of positive roots. It follows easily that

$$|\lambda_\pi|^2 \leq \kappa_\pi \leq |\lambda_\pi|^2 + 2|\lambda_\pi| |\rho| \leq C(1 + |\lambda_\pi|^2) \quad (3.11)$$

where $C > 1$ is a constant.

We also need the fact (which is implicit in the proof of Lemma 1.3 in [37]) that there exists $C_1, C_2 > 0$ such that for all $\lambda \in P \cap D$, there exists $n = (n_1, \dots, n_r) \in \mathbb{Z}^r$ such that

$$C_1 \|n\| \leq |\lambda| \leq C_2 \|n\|, \quad (3.12)$$

where $\|n\|^2 := n_1^2 + \dots + n_r^2$.

The final result we need from Sugiura [37] is Lemma 1.3 therein that

$$\zeta(s) := \sum_{\lambda \in P \cap D - \{0\}} \langle \lambda, \lambda \rangle^{-s} \quad (3.13)$$

converges if $2s > r$.

3.2 Sobolev Spaces

Let $\{X_1, \dots, X_d\}$ be a basis for the Lie algebra \mathfrak{g} of left-invariant vector fields. We define the Sobolev space $\mathcal{H}_p(G)$ by the prescription

$$\mathcal{H}_p(G) := \{f \in L^2(G); X_{i_1} \cdots X_{i_k} f \in L^2(G); 1 \leq k \leq p, i_1, \dots, i_k = 1, \dots, d\}.$$

It is a complex separable Hilbert space with associated norm

$$\|f\|_p^2 = \|f\|^2 + \sum_{i_1, \dots, i_k} \|X_{i_1} \cdots X_{i_k} f\|^2.$$

It is not difficult to show that an equivalent norm is given by

$$\|f\|_p^2 = \sum_{\pi \in \widehat{G}} d_\pi (1 + \kappa_\pi)^p \text{tr}(\widehat{f}(\pi) \widehat{f}(\pi)^*), \quad (3.14)$$

where $\widehat{f}(\pi) := \int_G \pi(\sigma^{-1}) f(\sigma) d\sigma$ is the Fourier transform¹ (and we are abusing notation by using $\|\cdot\|$ in each case.)

As is pointed out in [33], section 10.3.1, $\mathcal{H}_p(G)$ coincides with the usual Sobolev space on a manifold constructed using partitions of unity. So in particular the Sobolev embedding theorem extends to this context and hence

$$C^\infty(G) \supseteq \bigcap_{k \in \mathbb{N}} \mathcal{H}_k(G).$$

¹Note that we are here using the analyst's convention for Fourier transforms of functions which, as usual, is not quite consistent with the probabilist's convention for Fourier transforms of measures.

3.3 A Regularity Result

We summarise the results we need on regularity in the following:

Proposition 3.1 *Let $\mu \in \mathcal{M}_c(G)$ with $\widehat{\mu}(\pi) = c_\pi I_\pi$ for all $\pi \in \widehat{G}$.*

1. *The measure μ has a square-integrable density if and only if*

$$\sum_{\pi \in \widehat{G}} d_\pi^2 |c_\pi|^2 < \infty \quad (3.15)$$

2. *The measure μ has a continuous density if*

$$\sum_{\pi \in \widehat{G}} d_\pi^2 |c_\pi| < \infty, \quad (3.16)$$

3. *The measure μ has a C^k density if*

$$\sum_{\pi \in \widehat{G}} d_\pi^2 (1 + \kappa_\pi)^p |c_\pi|^2 < \infty. \quad (3.17)$$

where $p > k + \frac{d}{2}$.

Proof. (1) follows from Theorem 3.1 in [3] and (2) from Proposition 6.6.1 in [11]. (3) is a straightforward consequence of the Sobolev embedding theorem. \square

3.4 Examples

Now we consider different families of measures and apply Proposition 3.1. In all cases we take $\mu \in \text{CID}_{\mathbb{R}}(G)$ so that $c_\pi = e^{-\eta(\kappa_\pi^{\frac{1}{2}})}$.

3.4.1 The Case where there is a non-trivial Gaussian Component

We say that μ has a *non-trivial Gaussian component* if η is such that $\sigma > 0$ in (2.4). We can obtain many examples of such measures by defining $\mu = \mu_1 * \mu_2$ where μ_1 is Gaussian and μ_2 is of compound Poisson type or is obtained by subordination as in section 2.

We show that μ has a C^∞ -density for all $\sigma > 0$. To prove this we use (3.17), (3.8), (3.11) and (3.12) and the fact that $\eta(u) \geq \frac{1}{2}\sigma^2 u^2$ for all $u \in \mathbb{R}$

to see that for all $k \in \mathbb{N}$

$$\begin{aligned}
\sum_{\pi \in \widehat{G}} d_{\pi}^2 (1 + \kappa_{\pi})^k c_{\pi}^2 &\leq \sum_{\pi \in \widehat{G}} d_{\pi}^2 (1 + \kappa_{\pi})^k \exp \{-\sigma^2 \kappa_{\pi}\} \\
&\leq M \sum_{\lambda \in P \cap D} |\lambda|^{2m} (1 + |\lambda|^2)^k \exp \{-\sigma^2 |\lambda|^2\} \\
&\leq K_1 \sum_{n \in \mathbb{Z}^r} \|n\|^{2m} (1 + \|n\|^2)^k \exp \{-K_2 \|n\|^2\} \\
&= K_1 \sum_{j=0}^{\infty} a(j) j^m (1 + j)^k \exp \{-K_2 j\} \\
&\leq K_1 \sum_{j=0}^{\infty} j^m (2\sqrt{j} + 1)^r (1 + j)^k \exp \{-K_2 j\} < \infty
\end{aligned}$$

where $M, K_1, K_2 > 0$, $a(j) := \#\{n \in \mathbb{Z}^r; \|n\|^2 = j\}$ and we use the fact that $a(j) \leq (2\sqrt{j} + 1)^r$ for all $j \in \mathbb{N}$.²

3.4.2 Stable-type densities

Take $c_{\pi} = \exp \left\{ -b^{\alpha} \kappa_{\pi}^{\frac{\alpha}{2}} \right\}$ with $0 < \alpha < 2$. Again we show that the densities are C^{∞} . Indeed arguing as above we have

$$\begin{aligned}
\sum_{\pi \in \widehat{G}} d_{\pi}^2 (1 + \kappa_{\pi})^k c_{\pi}^2 &\leq M \sum_{\lambda \in P \cap D} |\lambda|^{2m} (1 + |\lambda|^2)^k \exp \{-2b^{\alpha} |\lambda|^{\alpha}\} \\
&\leq K_3 \sum_{j=0}^{\infty} j^m (2\sqrt{j} + 1)^r (1 + j)^k \exp \left\{ -K_4 j^{\frac{\alpha}{2}} \right\},
\end{aligned}$$

where $K_3, K_4 > 0$. To see that the series converges, it is sufficient to show that $\sum_{n=1}^{\infty} n^{\kappa} e^{-n^{\beta}}$ converges for all $\kappa \geq 0$ where $0 < \beta < 1$. This follows by comparison with $\sum_{n=1}^{\infty} \frac{1}{n^2}$ since $\lim_{n \rightarrow \infty} n^{\kappa+2} e^{-n^{\beta}} = \lim_{x \rightarrow \infty} x^{\frac{\kappa+2}{\beta}} e^{-x} = 0$.

3.4.3 Relativistic Schrödinger density

Here we have

²Of course in the pure Gaussian case, smoothness of the density is well-known and can be proved using pde techniques.

$$\begin{aligned}
\sum_{\pi \in \widehat{G}} d_{\pi}^2 (1 + \kappa_{\pi})^k c_{\pi}^2 &\leq e^{2m} \sum_{\lambda \in P \cap D} |\lambda|^{2m} |1 + |\lambda|^2|^k e^{-2\sqrt{m^2 + |\lambda|^2}} \\
&\leq e^{2m} \sum_{\lambda \in P \cap D} |\lambda|^{2m} |1 + |\lambda|^2|^k e^{-2|\lambda|} < \infty,
\end{aligned}$$

so by the result of (3.4.2) (with $\alpha = 1$) this case also yields a C^{∞} density.

3.4.4 Laplace density

In this case we take $c_{\pi} = (1 + \beta^2 \kappa_{\pi})^{-1}$. We restrict ourselves to seeking an L^2 -density. Applying (3.15) we use (3.8), (3.11) and (3.13) to obtain

$$\sum_{\pi \in \widehat{G - \{\delta\}}} \frac{d_{\pi}^2}{(1 + \beta^2 \kappa_{\pi})^2} \leq \frac{N}{\beta^2} \sum_{\lambda \in P \cap D - \{0\}} \frac{|\lambda|^{2m}}{|\lambda|^4} = \frac{N}{\beta^2} \zeta(2 - m),$$

where $N > 0$.

By Sugiura's convergence result for $\zeta(s)$ we see that a sufficient condition for convergence is $m < 2 - \frac{r}{2}$. Hence by (3.9), $\dim(G) \in \{1, 2, 3\}$. So for example, the Laplace distribution has a square-integrable density on the groups $\text{SO}(3)$, $\text{SU}(2)$ and $\text{Sp}(1)$, each of which has dimension 3 and rank 1.

4 Pseudo-differential Operator Representations

In this section G is an arbitrary compact group. Let $(\mu_t, t \geq 0)$ be a vaguely continuous convolution semigroup of probability measures on G wherein $\mu_0 = \delta_e$. It then follows that μ_t is infinitely divisible for each $t \geq 0$. We let $(T_t, t \geq 0)$ be the associated C_0 semigroup on $C(G)$ (*Hunt semigroup*) defined by

$$T_t f(\sigma) = \int_G f(\sigma\tau) \mu_t(d\tau),$$

for all $t \geq 0$. Necessary and sufficient conditions for a densely defined linear operator to extend to the infinitesimal generator of $(T_t, t \geq 0)$ were found by Hunt [20] (see [29] for a modern treatment) in the case of a Lie group and generalised by Born [9] to locally compact groups.

$(T_t, t \geq 0)$ extends to a positivity-preserving contraction semigroup on $L^2(G) := L^2(G, \mathbb{C})$ and from now on we will always work with this extended action. Our aim in this section is to represent the semigroup and its generator

as pseudo-differential operators using Peter-Weyl theory (c.f. [33]). If $A \in M_n(\mathbb{C})$, we define its Hilbert-Schmidt norm by $\|A\|_{\text{HS}} := \text{tr}(AA^*)^{\frac{1}{2}}$.

The celebrated Peter-Weyl theorem asserts that $f \in L^2(G)$ has the associated Fourier series $\sum_{\pi \in \widehat{G}} d_\pi \text{tr}(\widehat{f}(\pi)\pi)$ and we will make frequent use of Plancherel's theorem in this context which tells us that

$$\|f\|^2 = \sum_{\pi \in \widehat{G}} d_\pi \|\widehat{f}(\pi)\|_{\text{HS}}^2.$$

We also need the corresponding Parseval identity:

$$\langle f, g \rangle = \sum_{\pi \in \widehat{G}} d_\pi \text{tr}(\widehat{f}(\pi)\widehat{g}(\pi)^*),$$

for $f, g \in L^2(G)$.

We say that a densely defined linear operator S on $L^2(G)$ has a (*simple*) *pseudo-differential operator representation* if for each $\pi \in \widehat{G}$ there exists a $d_\pi \times d_\pi$ matrix $\sigma_S(\pi)$ such that

$$\widehat{Sf}(\pi) = \sigma_S(\pi)\widehat{f}(\pi),$$

for all $f \in \text{Dom}(S)$ and all $\pi \in \widehat{G}$.

We call σ_S the *symbol* of the operator S . The word “simple” is included as we don't require the symbol be a function defined on $G \times \widehat{G}$ as in [33]. Indeed such a more complicated class of symbols will be associated with representations of more general Feller-Markov semigroups (see [23] for the case where G is the real numbers.)

Theorem 4.1 *For each $t \geq 0$, T_t is a pseudo-differential operator with symbol $\widehat{\mu}_t(\pi)$ at $\pi \in \widehat{G}$.*

Proof. For each $\rho \in G$ let R_ρ denote right translation so that $R_\rho f(\sigma) = f(\sigma\rho)$ for each $f \in L^2(G), \sigma \in G$. We will need the fact that $\widehat{R_\rho f}(\pi) = \pi(\rho)\widehat{f}(\pi)$ for each $\pi \in \widehat{G}$.

By Fubini's theorem and the Parseval identity,

$$\begin{aligned} \|T_t f\|^2 &= \int_G \int_G \int_G f(\sigma\tau) \overline{f(\sigma\rho)} \mu_t(d\tau) \mu_t(d\rho) d\sigma \\ &= \int_G \int_G \int_G R_\tau f(\sigma) \overline{R_\rho f(\sigma)} d\sigma \mu_t(d\tau) \mu_t(d\rho) \\ &= \int_G \int_G \sum_{\pi \in \widehat{G}} d_\pi \text{tr}(\pi(\tau)\widehat{f}(\pi)\widehat{f}(\pi)^*\pi(\rho)^*) \mu_t(d\tau) \mu_t(d\rho). \end{aligned}$$

We can use Fubini's theorem to interchange summation and integration since by the contraction property of T_t ,

$$\int_G \int_G \sum_{\pi \in \widehat{G}} d_\pi \text{tr}(\pi(\tau) \widehat{f}(\pi) \widehat{f}(\pi)^* \pi(\rho)^*) \mu_t(d\tau) \mu_t(d\rho) \leq \|f\|^2.$$

Hence we have

$$\begin{aligned} \|T_t f\|^2 &= \sum_{\pi \in \widehat{G}} d_\pi \text{tr} \left(\int_G \pi(\tau) \mu_t(d\tau) \widehat{f}(\pi) \widehat{f}(\pi)^* \int_G \pi(\rho)^* \mu_t(d\rho) \right) \\ &= \sum_{\pi \in \widehat{G}} d_\pi \|\widehat{\mu}_t(\pi) \widehat{f}(\pi)\|_{hs}^2, \end{aligned}$$

and the result follows. \square

Let \mathcal{A} be the infinitesimal generator of $(T_t, t \geq 0)$. We here use the fact that for each $t \geq 0, \pi \in \widehat{G}, \widehat{\mu}_t(\pi) = e^{t\mathcal{L}_\pi}$ where \mathcal{L}_π is a $d_\pi \times d_\pi$ matrix (see [30], [18], [1] where an explicit ‘‘Lévy-Khintchine type’’ representation of \mathcal{L}_π can be found when G is a Lie group.)

Theorem 4.2 \mathcal{A} is a pseudo-differential operator with symbol \mathcal{L}_π at $\pi \in \widehat{G}$.

Proof. For each $f \in \text{Dom}(\mathcal{A}), g \in L^2(G)$, we have by Parseval's identity and Theorem 4.1

$$\begin{aligned} \langle \mathcal{A}f, g \rangle &= \lim_{t \rightarrow 0} \sum_{\pi \in \widehat{G}} d_\pi \text{tr} \left(\frac{(\widehat{\mu}_t(\pi) - I_\pi)}{t} \widehat{f}(\pi) \widehat{g}(\pi)^* \right) \\ &= \lim_{t \rightarrow 0} \sum_{\pi \in \widehat{G}} d_\pi \text{tr} \left(\frac{(e^{t\mathcal{L}_\pi} - I_\pi)}{t} \widehat{f}(\pi) \widehat{g}(\pi)^* \right) \end{aligned}$$

Now fix $\pi' \in \widehat{G}$ and let $g \in \mathcal{M}_{\pi'}$ where $\mathcal{M}_{\pi'}$ is the subspace of $L^2(G)$ generated by mappings of the form $\sigma \rightarrow \langle \pi'(\sigma)u, v \rangle$ for $u, v \in V_{\pi'}$. It follows from the Peter-Weyl theorem that $\widehat{g}(\pi) = 0$ if $\pi \neq \pi'$ and so

$$\begin{aligned} \langle \mathcal{A}f, g \rangle &= d_{\pi'} \lim_{t \rightarrow 0} \text{tr} \left(\frac{(e^{t\mathcal{L}_{\pi'}} - I_{\pi'})}{t} \widehat{f}(\pi') \widehat{g}(\pi')^* \right) \\ &= d_{\pi'} \text{tr}(\mathcal{L}_{\pi'} \widehat{f}(\pi') \widehat{g}(\pi')^*) \\ &= \sum_{\pi \in \widehat{G}} d_\pi \text{tr}(\mathcal{L}_\pi \widehat{f}(\pi) \widehat{g}(\pi)^*). \end{aligned}$$

The required result follows from the Parseval identity since (by the Peter-Weyl theorem) $L^2(G)$ is the closure of $\bigoplus_{\pi \in \widehat{G}} \mathcal{M}_\pi$ (see e.g. [11], p.108). \square

For completeness we will also give the pseudo-differential operator representation of the resolvent $R_\lambda := (\lambda I - \mathcal{A})^{-1}$ for $\lambda > 0$.

Proposition 4.1 *For each $\lambda > 0$, R_λ is a pseudo-differential operator having symbol $(\lambda I_\pi - \mathcal{L}_\pi)^{-1}$ at $\pi \in \widehat{G}$.*

Proof. First note that $(\lambda I_\pi - \mathcal{L}_\pi)^{-1}$ always exists since the eigenvalues of the matrix \mathcal{L}_π have non-positive real parts. We use the fact that for all $\lambda > 0$, $R_\lambda = \int_0^\infty e^{-\lambda t} T_t dt$. Then for all $f, g \in L^2(G)$, by Theorem 4.1

$$\langle R_\lambda f, g \rangle = \sum_{\pi \in \widehat{G}} d_\pi \int_0^\infty e^{-\lambda t} \text{tr}(e^{t\mathcal{L}_\pi} \widehat{f}(\pi) \widehat{g}(\pi)^*) dt.$$

The result follows from Fubini's theorem using the estimate

$$\sum_{\pi \in \widehat{G}} d_\pi \int_0^\infty e^{-\lambda t} |\text{tr}(e^{t\mathcal{L}_\pi} \widehat{f}(\pi) \widehat{g}(\pi)^*)| dt \leq \frac{1}{\lambda} \|f\| \cdot \|g\|,$$

which is obtained by routine computations. \square

Now we assume that $\mu_1 \in \text{CID}_{\mathbb{R}}(G)$. It follows that $\mu_t \in \text{CID}_{\mathbb{R}}(G)$ for all $t \geq 0$ and that $\widehat{\mu}_t(\pi) = e^{-t\eta(\kappa_\pi^{\frac{1}{2}})} I_\pi$ for each $\pi \in \widehat{G}$ for some negative definite function η and \mathcal{A} has symbol whose value at $\pi \in \widehat{G}$ is $-\eta(\kappa_\pi^{\frac{1}{2}}) I_\pi$.

Theorem 4.3 *If G is a compact Lie group and $\mu_1 \in \text{CID}_{\mathbb{R}}(G)$ then for all $p \geq 2$, $\mathcal{H}_p(G) \subseteq \text{Dom}(\mathcal{A})$ and \mathcal{A} is a bounded linear operator from $\mathcal{H}_p(G)$ to $\mathcal{H}_{p-2}(G)$.*

Proof (c.f. Theorem 10.81 in [33] pp.571-2.) We will make use of the fact that there exists $K > 0$ such that $|\eta(u)| \leq K(1 + |u|^2)$ for all $u \in \mathbb{R}$ (see e.g. [4] p.31.) By Theorem 4.2 and (3.14) for each $f \in \mathcal{H}_p(G)$

$$\begin{aligned} \|\mathcal{A}f\|_{p-2}^2 &= \sum_{\pi \in \widehat{G}} d_\pi (1 + \kappa_\pi)^{p-2} \|\mathcal{L}_\pi \widehat{f}(\pi)\|_{\text{hs}}^2 \\ &= \sum_{\pi \in \widehat{G}} d_\pi (1 + \kappa_\pi)^{p-2} |\eta(\kappa_\pi^{\frac{1}{2}})|^2 \|\widehat{f}(\pi)\|_{\text{hs}}^2 \\ &\leq K \sum_{\pi \in \widehat{G}} d_\pi (1 + \kappa_\pi)^p \|\widehat{f}(\pi)\|_{\text{hs}}^2 \\ &= K \|f\|_p^2. \end{aligned}$$

In particular it follows that $\|\mathcal{A}f\|^2 < \infty$ and so $f \in \text{Dom}(\mathcal{A})$. \square

5 Transition Densities for Convolution Semigroups

In this section we continue to work with the Hunt semigroup $(T_t, t \geq 0)$ acting on the space $L^2(G)$ that is associated to the convolution semigroup of measures $(\mu_t, t \geq 0)$ on a compact group G . Let $L_c^2(G) := \{f \in L^2(G), f(g\sigma g^{-1}) = f(\sigma) \text{ for all } \sigma, g \in G\}$. It is well-known that $\{\chi_\pi, \pi \in \widehat{G}\}$ is a complete orthonormal basis for $L_c^2(G)$ (see e.g. [11] Proposition 6.5.3, p.117.)

Proposition 5.1 *If $\mu_t \in M_c(G)$ for some $t \geq 0$ then $T_t(L_c^2(G)) \subseteq L_c^2(G)$.*

Proof. For all $\sigma, g \in G, f \in L_c^2(G)$ we have

$$\begin{aligned} T_t f(g\sigma g^{-1}) &= \int_G f(g\sigma g^{-1}\tau) \mu_t(d\tau) \\ &= \int_G f(g\sigma g^{-1}\tau) \mu_t(dg^{-1}\tau g) \\ &= \int_G f(g\sigma\tau g^{-1}) \mu_t(d\tau) \\ &= T_t f(\sigma). \quad \square \end{aligned}$$

Now suppose that $\mu_t \in M_{cs}(G)$ for all $t \geq 0$. This implies in particular that $(T_t, t \geq 0)$ is self-adjoint in $L^2(G)$ and hence in $L_c^2(G)$ (see [5], [27]). By Proposition 2.1, we also have that there exists $\alpha_\pi \leq 0$ for each $\pi \in \widehat{G}$ such that $\widehat{\mu}_t(\pi) = e^{t\alpha_\pi} I_\pi$.

Theorem 5.1 *If $\mu_t \in M_{cs}(G)$ for all $t \geq 0$ then $\{\chi_\pi, \pi \in \widehat{G}\}$ is a complete set of eigenvectors for the action of T_t on $L_c^2(G)$ and*

$$T_t \chi_\pi = e^{t\alpha_\pi} \chi_\pi \quad (5.18)$$

for all $\pi \in \widehat{G}, t \geq 0$.

Proof. For all $\sigma \in G, t \geq 0$,

$$\begin{aligned} T_t \chi_\pi(\sigma) &= \int_G \chi_\pi(\sigma\tau) \mu_t(d\tau) \\ &= \int_G \text{tr}(\pi(\sigma)\pi(\tau)) \mu_t(d\tau) \\ &= \text{tr} \left(\pi(\sigma) \int_G \pi(\tau) \mu_t(d\tau) \right) \\ &= e^{t\alpha_\pi} \chi_\pi(\sigma). \quad \square \end{aligned}$$

It is shown in [5] that T_t is trace-class for $t > 0$ if and only if μ_t has a square-integrable density. In this case we have

$$\mathrm{tr}(T_t) = \sum_{\pi \in \widehat{G}} e^{t\alpha_\pi} \quad (5.19)$$

for $t > 0$, where tr denote the trace in the Hilbert space $L_c^2(G)$.

From now on we assume that for $t > 0$, $\mu_t \in \mathcal{M}_{cs}(G)$ has a density $k_t \in L_c^2(G, \mathbb{R})$. We define the *transition density* $h_t \in L^2(G \times G, \mathbb{R})$ by

$$h_t(\sigma, \tau) := k_t(\sigma^{-1}\tau),$$

for each $t > 0, \sigma, \tau \in G$. Indeed h_t is precisely the transition probability density at time t of a G -valued Lévy process whose law at time t is k_t .

Note that for each $\sigma \in G, (t, \rho) \rightarrow h_t(\sigma, \rho)$ satisfies the backward equation (in the distributional sense)

$$\frac{\partial h_t}{\partial t}(\sigma, \rho) = \mathcal{A}h_t(\sigma, \rho),$$

with $h_t(\sigma, \rho) \rightarrow \delta_\sigma(\rho)$ as $t \rightarrow 0$. For example, if $(\mu_t, t \geq 0)$ is the Brownian motion semigroup which is characterised by $\widehat{\mu}_t(\pi) = e^{-\frac{t}{2}\kappa_\pi} I_\pi$ for each $\pi \in \widehat{G}$, then h_t is the well-known heat kernel and for this reason we will sometimes refer to our more general h_t as the *transition kernel*.

Theorem 5.2 *For each $t > 0$,*

1.

$$\int_G h_t(g^{-1}\sigma, \rho g^{-1}) dg = \sum_{\pi \in \widehat{G}} e^{t\alpha_\pi} \overline{\chi_\pi(\sigma)} \chi_\pi(\rho), \quad (5.20)$$

for all $\sigma, \rho \in G$.

2.

$$\mathrm{tr}(T_t) = \int_G \int_G k_t(\rho^{-1}g\rho g^{-1}) dg d\rho \quad (5.21)$$

Proof.

1. By (3.7)

$$h_t(\sigma, \rho) = k_t(\sigma^{-1}\rho) = \sum_{\pi \in \widehat{G}} d_\pi e^{t\alpha_\pi} \chi_\pi(\sigma^{-1}\rho), \quad (5.22)$$

and so

$$\begin{aligned} \int_G h_t(g^{-1}\sigma, \rho g^{-1}) dg &= \sum_{\pi \in \widehat{G}} d_\pi e^{t\alpha_\pi} \int_G \chi_\pi(\sigma^{-1} g \rho g^{-1}) dg \\ &= \sum_{\pi \in \widehat{G}} e^{t\alpha_\pi} \overline{\chi_\pi(\sigma)} \chi_\pi(\rho), \end{aligned}$$

by Proposition 6.5.2 in [11] (p.116.) The interchange of integral and sum is justified by Fubini's theorem since

$$\sum_{\pi \in \widehat{G}} d_\pi e^{t\alpha_\pi} \int_G |\chi_\pi(\sigma^{-1} g \rho g^{-1})| dg \leq \sum_{\pi \in \widehat{G}} d_\pi^2 e^{t\alpha_\pi} < \infty$$

by (3.15) since $k_t \in L^2(G)$ for each $t > 0$. Here we have used the crude estimate $\sup_{\sigma \in G} |\chi_\pi(\sigma)| \leq d_\pi$.

2. Put $\rho = \sigma$ in (5.20) and then integrate both sides with respect to σ using the fact that $\int_G |\chi_\pi(\sigma)|^2 d\sigma = 1$. The result then follows from (5.19). Note that the interchange of integral and summation is justified by Fubini's theorem using a similar argument to that presented in (1). \square

Corollary 5.1 *If k_t is continuous for each $t > 0$*

$$k_t(e) = \sum_{\pi \in \widehat{G}} d_\pi^2 e^{t\alpha_\pi}. \quad (5.23)$$

Proof. Put $\sigma = \rho$ in (5.22) (or argue directly from (3.7).) \square

We now work in the Hilbert space $L^2(G)$ and we use Tr to denote the trace in this Hilbert space. By the Peter-Weyl theorem $\{d_\pi^{\frac{1}{2}} \pi_{ij}, 1 \leq i, j \leq d_\pi, \pi \in \widehat{G}\}$ is a complete orthonormal basis for $L^2(G)$ where π_{ij} denotes the co-ordinate function $\pi_{ij}(\sigma) := \pi(\sigma)_{ij}$, for each $\sigma \in G$.

The following two results are well known for the heat kernel (see e.g. Chapter 12 of [14]). Here we extend them to more general Hunt semigroups.

Theorem 5.3 *1. For each $t \geq 0$, the set $\{\pi_{ij}, 1 \leq i, j \leq d_\pi, \pi \in \widehat{G}\}$ is a complete orthogonal set of eigenvectors for T_t and*

$$T_t \pi_{ij} = e^{t\alpha_\pi} \pi_{ij} \quad (5.24)$$

for each $1 \leq i, j \leq d_\pi, \pi \in \widehat{G}$.

2. If k_t is continuous for each $t > 0$,

$$k_t(e) = \text{Tr}(T_t). \quad (5.25)$$

Proof.

1. For all $\sigma \in G$.

$$\begin{aligned} T_t \pi_{ij}(\sigma) &= \int_G \pi_{ij}(\sigma\tau) \mu_t(d\tau) \\ &= \sum_{k=1}^{d_\pi} \pi_{ik}(\sigma) \int_G \pi_{kj}(\tau) \mu_t(d\tau) \\ &= \sum_{k=1}^{d_\pi} \pi_{ik}(\sigma) \widehat{\mu}_t(\pi)_{kj} \\ &= e^{t\alpha_\pi} \pi_{ij}(\sigma). \end{aligned}$$

2. Since each eigenvalue $e^{t\alpha_\pi}$ has multiplicity d_π^2 on the closed subspace of $L^2(G)$ spanned by $\{\pi_{ij}, 1 \leq i, j \leq d_\pi\}$ it is clear that

$$\text{Tr}(T_t) = \sum_{\pi \in \widehat{G}} d_\pi^2 e^{t\alpha_\pi},$$

and the result follows from (5.23). \square

The next result can be deduced directly from (3.7). We give a direct proof to make the paper more self-contained. Note that results of this type are well-known for Markov processes taking values in compact metric spaces and having suitably square-integrable transition probabilities (see Theorem 6.4 in [15].)

Theorem 5.4 *If k_t is continuous for each $t > 0$*

$$h_t(\sigma, \rho) = \sum_{\pi \in \widehat{G}} \sum_{i,j=1}^{d_\pi} d_\pi e^{t\alpha_\pi} \overline{\pi_{ij}(\sigma)} \pi_{ij}(\rho), \quad (5.26)$$

for all $\sigma, \rho \in G$.

Proof. For each $\sigma \in G$ let L_σ denote left translation so that $L_\sigma f(\rho) = f(\sigma^{-1}\rho)$ for each $f \in L^2(G), \rho \in G$. By Fourier expansion and using (5.24)

$$\begin{aligned}
L_\sigma k_t &= \sum_{\pi \in \widehat{G}} \sum_{i,j=1}^{d_\pi} d_\pi \langle L_\sigma k_t, \pi_{ij} \rangle \pi_{ij} \\
&= \sum_{\pi \in \widehat{G}} \sum_{i,j=1}^{d_\pi} d_\pi \left(\int_G k_t(\sigma^{-1}\tau) \overline{\pi_{ij}(\tau)} d\tau \right) \pi_{ij} \\
&= \sum_{\pi \in \widehat{G}} \sum_{i,j=1}^{d_\pi} d_\pi \left(\int_G k_t(\tau) \overline{\pi_{ij}(\sigma\tau)} d\tau \right) \pi_{ij} \\
&= \sum_{\pi \in \widehat{G}} \sum_{i,j=1}^{d_\pi} d_\pi T_t \overline{\pi_{ij}(\sigma)} \pi_{ij} \\
&= \sum_{\pi \in \widehat{G}} \sum_{i,j=1}^{d_\pi} d_\pi e^{t\alpha_\pi} \overline{\pi_{ij}(\sigma)} \pi_{ij}.
\end{aligned}$$

Since $\sup_{\sigma, \rho \in G} \left| \sum_{i,j=1}^{d_\pi} \overline{\pi_{ij}(\sigma)} \pi_{ij}(\rho) \right| = \sup_{\sigma, \rho \in G} |\text{tr}(\pi(\sigma^{-1}\rho))| \leq d_\pi$ we deduce uniform convergence of the series from (3.15) and so for all $\rho \in G$

$$h_t(\sigma, \rho) = L_\sigma k_t(\rho) = \sum_{\pi \in \widehat{G}} \sum_{i,j=1}^{d_\pi} d_\pi e^{t\alpha_\pi} \overline{\pi_{ij}(\sigma)} \pi_{ij}(\rho).$$

□

Corollary 5.2 *If k_t is continuous for each $t > 0$*

$$k_t(e) \geq \int_G \int_G k_t(\rho^{-1}g\rho g^{-1}) dg d\rho$$

with equality if and only if G is abelian.

Proof. The inequality follows from (5.19), (5.21), (5.23) and (5.25). If G is abelian, then equality is obvious. If G is non-abelian we must have $d_\pi > 1$ for at least one $\pi \in \widehat{G}$ and then it is clear that strict inequality holds. □

6 Small Time Asymptotics of Densities

Assume that G is a compact semisimple Lie group. We would like to obtain an asymptotic expansion for $k_t(e)$ as $t \rightarrow 0$. We assume that $\mu_t \in \text{CID}_{\mathbb{R}}(G)$

for each $t \geq 0$. In this case if k_t is continuous for $t > 0$, we may follow the arguments on p.106 of [14] to obtain

$$\begin{aligned} k_t(e) = \text{Tr}(T_t) &= \sum_{\lambda \in P \cap D} d_{\lambda+\rho}^2 \exp\{-t\eta((|\lambda + \rho|^2 - |\rho|^2)^{\frac{1}{2}})\} \quad (6.27) \\ &= \sum_{\lambda \in P \cap D} d_{\lambda}^2 \exp\{-t\eta((|\lambda|^2 - |\rho|^2)^{\frac{1}{2}})\} \\ &= \frac{1}{|W|} \sum_{\lambda \in P} d_{\lambda}^2 \exp\{-t\eta((|\lambda|^2 - |\rho|^2)^{\frac{1}{2}})\}, \end{aligned}$$

where d_{λ} denotes the dimension of the representation space with highest weight λ and $|W|$ is the order of the Weyl group of G .

When $\eta(u) = \frac{|u|^2}{2}$, k_t is the density generating the heat kernel and it is known that as $t \rightarrow 0$,

$$k_t(e) \sim Ct^{-\frac{\dim(G)}{2}} e^{t|\rho|^2} \quad (6.28)$$

(see e.g. [14], p.109)) where $C > 0$.

We will examine the case where η is the characteristic exponent of a symmetric Cauchy distribution so that $\eta(u) = \sigma|u|$ for all $u \in \mathbb{R}$, where $\sigma > 0$.

Example 1: $G = \Pi^d$ where $\Pi := \mathbb{R}/\mathbb{Z}$

In this case $\widehat{G} = \mathbb{Z}^d$, each $d_{\pi} = 1$ and for each $\pi = n \in \mathbb{Z}^d$, $\kappa_{\pi} = 4\pi^2|n|^2$ where $n^2 = n_1^2 + \dots + n_d^2$ for $n = (n_1, \dots, n_d)$. The equation (6.27) then takes the form

$$k_t(e) = \sum_{n \in \mathbb{Z}^d} e^{-2\pi t \sigma |n|}.$$

When $d = 1$ we easily calculate

$$\begin{aligned} k_t(e) &= 1 + 2 \sum_{n=1}^{\infty} e^{-2\pi t \sigma n} \\ &= \coth(\pi \sigma t) \sim \frac{1}{\pi \sigma t}, \text{ as } t \rightarrow 0. \end{aligned}$$

When $d > 1$, we apply the Poisson summation formula to obtain

$$\begin{aligned} k_t(e) &= \frac{\Gamma(\frac{d+1}{2})}{\pi^{\frac{d+1}{2}}} \sum_{m \in \mathbb{Z}^d} \frac{\sigma t}{(\sigma^2 t^2 + |m|^2)^{\frac{d+1}{2}}} \\ &\sim \frac{\Gamma(\frac{d+1}{2})}{\sigma^d \pi^{\frac{d+1}{2}}} \frac{1}{t^d}, \text{ as } t \rightarrow 0. \end{aligned}$$

Example 2: $G = SU(2)$. In this case $\widehat{G} \cong \mathbb{Z}_+$ with $d_n = n + 1$ and for each $n \in \mathbb{Z}_+$, $\kappa_n = n(n + 2)$. Hence

$$\begin{aligned} k_t(e) &= \sum_{n=0}^{\infty} (n+1)^2 e^{-t\sigma\sqrt{n(n+2)}} \\ &= \sum_{n=0}^{\infty} (n+1)^2 e^{-t\sigma\sqrt{(n+1)^2-1}} \\ &= \sum_{m=1}^{\infty} m^2 e^{-t\sigma\sqrt{m^2-1}} \end{aligned}$$

From this we get the easy estimate

$$e^{-t\sigma} \sum_{m=1}^{\infty} m^2 e^{-t\sigma m} \leq k_t(e) \leq e^{t\sigma} \sum_{m=1}^{\infty} m^2 e^{-t\sigma m}. \quad (6.29)$$

Now define $g(t) := \sum_{m=1}^{\infty} e^{-t\sigma m}$ for $t \in (0, \infty)$. The function g is C^∞ and we have

$$\begin{aligned} \sum_{m=1}^{\infty} m^2 e^{-t\sigma m} &= \frac{1}{\sigma^2} \frac{d^2}{dt^2} g(t) \\ &= \frac{1}{\sigma^2} \frac{d^2}{dt^2} \left(\frac{e^{-t\sigma}}{1 - e^{-t\sigma}} \right) \\ &= \frac{e^{-t\sigma}}{(1 - e^{-t\sigma})^2} \coth \left(\frac{\sigma t}{2} \right), \end{aligned}$$

and hence we conclude that

$$k_t(e) \sim \frac{2}{\sigma^3 t^3}, \text{ as } t \rightarrow 0.$$

This should be compared with the usual heat kernel where the following very precise asymptotic expansion is known (see [13], Proposition 2.3 p.662):

$$k_t(e) \sim 32\sqrt{2}\pi^2(4\pi\sigma t)^{-\frac{3}{2}} e^{\frac{\sigma t}{8}},$$

so the leading term has the slower decay

$$k_t(e) \sim 32\sqrt{2}\pi^2(4\pi\sigma)^{-\frac{3}{2}} t^{-\frac{3}{2}}.$$

It is not difficult to verify that the relativistic Schrödinger semigroup on $SU(2)$ with mass parameter m , for which $\eta(u) = (m^2 + u^2)^{\frac{1}{2}} - m$ for $u \in \mathbb{R}$, has exactly the same short time asymptotics as the Cauchy semigroup.

Example 3: $G = SO(3)$. Here we have $\widehat{G} \cong \mathbb{Z}_+$ with $d_n = 2n + 1$ and for each $n \in \mathbb{Z}_+$, $\kappa_n = n(n + 1)$. So we obtain

$$\begin{aligned} k_t(e) &= \sum_{n=0}^{\infty} (2n + 1)^2 e^{-t\sigma\sqrt{n(n+1)}} \\ &= \sum_{m \text{ odd}} m^2 e^{-t\frac{\sigma}{2}\sqrt{m^2-1}} \text{ as } t \rightarrow 0. \end{aligned}$$

Using the results obtained in Example 2 we find that

$$k_t(e) \sim \frac{8}{\sigma^3 t^3}, \text{ as } t \rightarrow 0.$$

Based on these calculations we conjecture that $k_t(e) \sim Ct^{-d}$ for the Cauchy process on an arbitrary compact semisimple Lie group of dimension d . It is also tempting to further conjecture that if k_t is associated to an arbitrary α -stable process so that $\eta(u) = |u|^\alpha$ where $0 < \alpha \leq 2$, then $k_t(e) \sim Ct^{-\frac{d}{\alpha}}$ as $t \rightarrow 0$. This is consistent with the known behaviour of densities of symmetric stable processes in Euclidean space (see e.g. [8]) where it essentially follows by scaling arguments. We remind the reader that this technology is not available on compact groups (see e.g. Theorem 2.2. in [2], p.117.)

Now let $N(\lambda)$ denote the number of eigenvalues of $-\mathcal{A}$ that do not exceed λ and note that for all $t > 0$,

$$\text{Tr}(T_t) = \int_0^\infty e^{-t\lambda} dN(\lambda).$$

If the above conjecture holds then by Karamata's Tauberian theorem we have

$$N(\lambda) \sim \frac{C\lambda^{\frac{d}{\alpha}}}{\Gamma(1 + \frac{d}{\alpha})}, \text{ as } \lambda \rightarrow \infty,$$

(c.f. Theorem 2.3 in [7].)

So far we know that this eigenvalue asymptotics is valid when $\alpha = 2$, and when $\alpha = 1$ with $G = \Pi^d$, $G = SU(2)$ and $G = SO(3)$.

References

- [1] D.Applebaum, Operator-valued stochastic differential equations arising from unitary group representations, *Journal of Theoretical Probability* **14**, 61-76 (2001)
- [2] D.Applebaum, Lévy processes in stochastic differential geometry in *Lévy Processes: Theory and Applications* ed. O.Barndorff-Nielsen, T.Mikosch, S.Resnick (Birkhäuser Boston Basel Berlin), 111-39 (2001)
- [3] D.Applebaum, Probability measures on compact groups which have square-integrable densities, *Bull. Lond. Math. Sci.* **40** 1038-44 (2008)
- [4] D.Applebaum *Lévy Processes and Stochastic Calculus* (second edition), Cambridge University Press (2009)
- [5] D.Applebaum, Some L^2 properties of semigroups of measures on Lie groups, *Semigroup Forum* **79**, 217-28 (2009)
- [6] C. Berg, G. Forst, *Potential Theory on Locally Compact Abelian Groups*, Springer-Verlag (1975).
- [7] R.M.Blumenthal, R.K.Gettoor, The asymptotic distribution of the eigenvalues for a class of Markov operators, *Pacific J. Math.* **9**, 399-408 (1959)
- [8] R.M.Blumenthal, R.K.Gettoor, Some theorems on stable processes, *Trans. Amer. Math. Soc.* **65**, 263-73 (1960)
- [9] E.Born, An explicit Lévy-Hinčín formula for convolution semigroups on locally compact groups, *J.Theor. Prob.* **2** 325-42 (1989)
- [10] K.D.Elworthy, Geometric aspects of diffusions on manifolds in *École d'Été de Probabilités de Saint-Flour XV-XVII, 1985-87*, 277-425, Lecture Notes in Math, **1362**, Springer, Berlin (1988)
- [11] J.Faraut, *Analysis on Lie Groups*, Cambridge University Press (2008)
- [12] H.D.Fegan, The heat equation and modular forms, *J.Differential Geometry* **13**, 589-602 (1978)
- [13] H.D.Fegan, The fundamental solution of the heat equation on a compact Lie group, *J.Differential Geometry* **18**, 659-668 (1983)
- [14] H.D.Fegan, *Introduction to Compact Lie Groups*, World Scientific (1991)

- [15] R.K.Gettoor, Markov operators and their associated semigroups, *Pacific J. Math.* **9**, 449-72 (1959)
- [16] K.E.Hare, The size of characters of compact Lie groups, *Studia Math.* **129** 1-18 (1998)
- [17] H.Heyer, L'analyse de Fourier non-commutative et applications à la théorie des probabilités, *Ann. Inst. Henri Poincaré (Prob. Stat.)* **4**(1968), 143-68
- [18] H.Heyer, Infinitely divisible probability measures on compact groups, in Lectures on Operator Algebras pp.55-249, Lecture Notes in Math. vol **247** Springer Berlin, Heidelberg, New York (1972)
- [19] H.Heyer, *Probability Measures on Locally Compact Groups*, Springer-Verlag, Berlin-Heidelberg (1977)
- [20] G.A.Hunt, Semigroups of measures on Lie groups, *Trans. Amer. Math. Soc.* **81**, 264-93 (1956)
- [21] T.Ichinose, Essential selfadjointness of the Weyl quantized relativistic Hamiltonian, *Ann. Inst. Henri Poincaré (Phys. Theor.)* **51**, 265-98 (1989)
- [22] N. Jacob, *Pseudo-Differential Operators and Markov Processes*, Akademie-Verlag, Mathematical Research 94 (1996).
- [23] N. Jacob, *Pseudo-Differential Operators and Markov Processes: 3, Markov Processes and Applications*, World Scientific (2005).
- [24] P.T.Kim, D.S.Richards, Deconvolution density estimators on compact Lie groups, *Contemp. Math.* **287**, 155-71 (2001)
- [25] A.W.Knapp, *Lie Groups Beyond an Introduction (second edition)*, Birkhäuser (1996, 2002)
- [26] J-Y Koo, P.T.Kim, Asymptotic minimax bounds for stochastic deconvolution over groups, *IEEE Trans. Inf. Theory* **54**, 289-98 (2008)
- [27] H.Kunita, Analyticity and injectivity of convolution semigroups on Lie groups, *J.Funct. Anal.* **165**, 80-100 (1999)
- [28] M.Liao, Lévy processes and Fourier analysis on compact Lie groups, *Ann. Prob.* **32** (2004), 1553-73

- [29] M.Liao, *Lévy Processes in Lie Groups*, Cambridge University Press (2004)
- [30] J.T-H.Lo, S-K.Ng, Characterizing Fourier series representations of probability distributions on compact Lie groups, *Siam J. Appl. Math.* **48** 222-8 (1988)
- [31] D.Ragozin, Central measures on compact simple Lie groups, *J.Funct. Anal.* **10**, 212-29 (1972)
- [32] S.Rosenberg, *The Laplacian on a Riemannian Manifold*, Cambridge University Press (1997)
- [33] M.Ruzhansky, V.Turunen, *Pseudo-differential Operators and Symmetries: Background Analysis and Advanced Topics*, Birkhäuser, Basel (2010)
- [34] S.Said, C.Lageman, N.LeBihan, J.H.Manton, Decomponding on compact Lie groups, *IEEE Trans. Inf. Theory* **56**, 2766-77 (2010)
- [35] K.-I.Sato, *Lévy Processes and Infinite Divisibility*, Cambridge University Press (1999)
- [36] E.Siebert, Fourier analysis and limit theorems for convolution semi-groups on a locally compact group, *Advances in Math.* **39** (1981), 111-54
- [37] M.Sugiura, Fourier series of smooth functions on compact Lie groups, *Osaka J.Math* **8**, 33-47 (1971)